

Stability of the Rational Homotopy Type of Moduli Spaces

Alexander A. Voronov

Dedicated to Jim Stasheff on the occasion of his sixtieth birthday.

ABSTRACT. We show that for $g \geq 2k + 3$ the k -rational homotopy type of the moduli space $\mathcal{M}_{g,n}$ of algebraic curves of genus g with n punctures is independent of g , and the space $\mathcal{M}_{g,n}$ is k -formal. This implies the existence of a limiting rational homotopy type of $\mathcal{M}_{g,n}$ as $g \rightarrow \infty$ and the formality of it.

Introduction

The description of the algebraic topology of the moduli space $\mathcal{M}_{g,n}$ of compact complex algebraic curves has long been a tantalizing problem. The idea of “stable cohomology” of $\mathcal{M}_{g,n}$ as the genus $g \rightarrow \infty$, brought in by J. L. Harer and D. Mumford, suggested a more graspable object to study as a first step. Mumford’s Conjecture [Mum83], stating that the stable cohomology is the polynomial algebra on the so-called Mumford-Morita-Miller classes, has become a new tantalizing problem since then. Recently M. Pikaart [Pik95] has proven that the Hodge structure on the stable cohomology is pure, thus providing more evidence to the conjecture, which easily implies purity.

Our purpose in this note is to show that the rational homotopy type of the moduli space $\mathcal{M}_{g,n}$ stabilizes as $g \rightarrow \infty$ and prove that the moduli space is formal in the stable limit. If the stable cohomology of the moduli space were just known to be the cohomology of a certain space $\mathcal{M}_{\infty,n}$, then a theorem of E. Miller and S. Morita [Mil86] asserting that $H^\bullet(\mathcal{M}_{\infty,n})$ is a free graded commutative algebra would immediately imply the formality of $\mathcal{M}_{\infty,n}$. Thus, the essential result of this paper is showing that the limit of the rational homotopy types of $\mathcal{M}_{g,n}$ as $g \rightarrow \infty$ exists. After this is done, the formality of the limit is automatic.

Here are few words about the structure of the paper. Sections 1 and 2 are dedicated to the description of the (nonstable) rational homotopy type of the moduli space $\mathcal{M}_{g,n}$. Section 3, where the stable rational homotopy type of $\mathcal{M}_{g,n}$ is studied, is independent of the first two sections.

It may be worth noting how this paper relates to Jim Stasheff’s work. Stasheff polyhedra K_n are known for many remarkable properties. One of them, that K_n is a

1991 *Mathematics Subject Classification*. Primary 14H10; Secondary 32G15, 55P62.
Research supported in part by an AMS Centennial Fellowship.

connected component of the real compactification of the moduli space of real projective lines with $n+1$ punctures, was pointed out by M. Kontsevich in [Kon94]. Thus, Stasheff polyhedra are real analogues of the compactified moduli spaces $\overline{\mathcal{M}}_{g,n}$. Or, perhaps, the moduli spaces $\overline{\mathcal{M}}_{g,n}$ are complex analogues of Stasheff polyhedra, the chicken and the egg problem. The topology of K_n is trivial (K_n is contractible), and the topology of $\mathcal{M}_{g,n}$ and $\overline{\mathcal{M}}_{g,n}$ is just the opposite. It is rather the combinatorics of Stasheff polyhedra which makes them very useful in the topology of loop spaces [Sta63]. The combinatorics of the genus zero spaces $\mathcal{M}_{0,n}$ has similarly proven to be useful in studying the topology of double loop spaces [GJ94] and the algebraic structure of 2d quantum field theory [KSV95, KSV96, KVZ97]. But what is the topology of these complex analogues of Stasheff polyhedra, especially for a high genus?

We also implicitly use a result of S. Halperin and Stasheff [HS79] throughout the paper: formality over any field of characteristic zero implies that over the rationals.

ACKNOWLEDGMENT. I am very grateful to Pierre Deligne, Dick Hain, Eduard Looijenga, and Martin Pikaart for helpful discussions. I would like to thank the Institute for Advanced Study in Princeton and the Research Institute for Mathematical Sciences in Kyoto for their hospitality during the summer 1997, when the essential part of the work on this paper was done.

1. The rational homotopy type of a complex smooth quasi-projective variety

Here we recall J. W. Morgan's description [Mor78] of a model of the rational homotopy type of the complement U of a normal crossings divisor $D = \bigcup_{i=1}^r D_i$ in a compact complex manifold X . The rational homotopy type will be understood in the sense of D. Sullivan [Sul77], see also P. A. Griffiths and Morgan [GM81] and D. Lehmann [Leh90]. It determines the usual rational homotopy type of a topological space if it is simply connected. In general it determines the rational nilpotent completion thereof.

The rational homotopy type of U is determined by the differential graded (DG) commutative algebra that is nothing but the first term $\mathcal{A}^\bullet = E_1^\bullet$ of the spectral sequence associated with the weight filtration on the log-complex $\Omega_X^{\bullet,\bullet}(\log D)$ of the smooth (p, q) -forms on U with logarithmic singularities along D . Otherwise, one can describe the same DG algebra as the second term of the Leray spectral sequence of inclusion $U \hookrightarrow X$. In any case, the model of U is given by the DG algebra $\mathcal{A}^\bullet$, where

$$\begin{aligned} \mathcal{A}^k &= \bigoplus_{p+q=k} \mathcal{A}^{p,q}, & \mathcal{A}^{p,q} &= \bigoplus_{\substack{S \subset \{1,2,\dots,r\} \\ |S|=-p}} H^{2p+q}(D_S, \mathbb{C}), \\ p \leq 0, \ q \geq 0, & & D_S &= \bigcap_{i \in S} D_i. \end{aligned}$$

The multiplication structure is given by

$$a \cdot a' = \begin{cases} (-1)^{pq'+\epsilon} (a|_{D_S \cap D_{S'}}) \cup (a'|_{D_S \cap D_{S'}}) & \text{if } S \cap S' = \emptyset, \\ 0 & \text{otherwise} \end{cases}$$

for $a \in H^{2p+q}(D_S, \mathbb{C})$ and $a' \in H^{2p'+q'}(D_{S'}, \mathbb{C})$, where ϵ is the sign of the shuffle putting the set $S \cap S'$ in an increasing order, assuming each of the subsets S and S' to be already in an increasing order. Note that this multiplication law makes $\mathcal{A}^{\bullet, \bullet}$ into a bigraded commutative algebra.

Finally, the differential $d : \mathcal{A}^{p,q} \rightarrow \mathcal{A}^{p+1,q}$ can be described as

$$da = \sum_{j=1}^{-p} (-1)^{j-1} (\iota_j)_* a,$$

where $a \in H^{2p+q}(D_S, \mathbb{C})$, $S = \{i_1, \dots, i_{-p}\}$, ι_j is the natural embedding $D_S \subset D_{S \setminus i_j}$, and $(\iota_j)_* : H^{2p+q}(D_S, \mathbb{C}) \rightarrow H^{2p+q+2}(D_{S \setminus i_j}, \mathbb{C})$ is the Gysin map.

P. Deligne [Del71] proved that the spectral sequence $E_1^{p,q} = \mathcal{A}^{p,q}$ degenerates at E_2 , that is, the cohomology of the DG algebra $\mathcal{A}^\bullet$ is equal to $H^\bullet(U, \mathbb{C})$. The very DG algebra $\mathcal{A}^\bullet$ describes the rational homotopy type of U , according to Morgan's theorem [Mor78].

2. The rational homotopy type of $\mathcal{M}_{g,n}$

Results of the previous section apply to the moduli space $\mathcal{M}_{g,n}$, which is the complement of a normal crossing divisor in the Deligne-Knudsen-Mumford compactification $\overline{\mathcal{M}}_{g,n}$. The problem that the space in question is a stack rather than variety does not arise, because we work with the **complex coefficients**, as we will assume throughout the paper. From the combinatorics of Deligne-Knudsen-Mumford's construction, we can say more specifically, cf. [KSV96], that

$$\mathcal{A}^{p,q} = \bigoplus_G \left(\bigotimes_{v \in G} H^\bullet(\overline{\mathcal{M}}_{g(v), n(v)}) \right)_{2p+q}^{\text{Aut}(G)},$$

the summation running over all stable labeled n -graphs G of genus $g(G) = g$ and $v(G) = -p + 1$ vertices. Here we refer to *graphs* of the following kind. Each graph is connected and has n *enumerated exterior edges*, edges which are incident with only one vertex of the graph. Each vertex v of the graph is *labeled* by a nonnegative integer $g(v)$, called the genus of a vertex. The *stability* condition means that any vertex v labeled by $g(v) = 1$ should be incident with at least one edge (i.e., be at least of valence one) and each vertex v with $g(v) = 0$ should be at least of valence three. The *genus* $g(G)$ of a graph G is given by the formula $g(G) = b_1(G) + \sum_v g(v)$, where $b_1(G)$ is the first Betti number of the graph. The *number* $n(v)$ is the valence of the vertex v and $\text{Aut}(G)$ is the *automorphism group of a graph* G (bijections on vertices and edges, preserving the exterior edges, the labels of vertices and the incidence relation). The subscript $2p + q$ in the formula refers to taking the homogeneous component of this degree. The differential $d : \mathcal{A}^{p,q} \rightarrow \mathcal{A}^{p+1,q}$ is induced by contracting interior edges in G , which corresponds to replacing a neighborhood of a double point on a curve by a cylinder.

Our goal here is to look at the stable (as $g \rightarrow \infty$) cohomology and rational homotopy type of the moduli space $\mathcal{M}_{g,n}$. The following result of Pikaart [Pik95] gives a certain clue on what is going on within a “stable range”.

THEOREM 2.1 (Pikaart). *The restriction mapping $H^k(\overline{\mathcal{M}}_{g,n}) \rightarrow H^k(\mathcal{M}_{g,n})$ is surjective for $k \leq (g - 1)/2$.*

COROLLARY. *For $p + q \leq (g - 1)/2$, the cohomology of $\mathcal{A}^{p,q}$ is nonzero only for $p = 0$, see Figure 1.*




FIGURE 1. The algebra $\mathcal{A}^{p,q}$. The shaded region is the “stable range”, where the cohomology is concentrated along the fat line.

PROOF. By our construction of the spectral sequence, the natural composite mapping $\mathcal{A}^{0,k} \rightarrow H^{0,k}(\mathcal{A}^{\bullet,\bullet}, d) \hookrightarrow H^k(\mathcal{A}^{\bullet,\bullet}, d)$ is the same as the restriction mapping $H^k(\overline{\mathcal{M}}_{g,n}) \rightarrow H^k(\mathcal{M}_{g,n})$. Therefore, the $p \neq 0$ columns of $\mathcal{A}^{p,q}$ do not contribute to the cohomology of the DG algebra $\mathcal{A}$ in the stable range. $\square$

3. The stable limit

The stable limit in cohomology of the moduli spaces $\mathcal{M}_{g,n}$ is achieved, roughly speaking, by gluing more and more handles to the complex curve. This yields an inductive system (of isomorphisms) on the level of cohomology. But since there is no natural mapping between the moduli spaces of different genera, one cannot speak of a limiting rational homotopy type. The question of taking the limit of the DG algebras $\mathcal{A}$ of Section 2 for $\mathcal{M}_{g,n}$ may not be so obviously resolved, either, because these algebras are constructed out of the cohomology of $\overline{\mathcal{M}}_{g',n}$, g' running between 0 and g . When $g \rightarrow \infty$, g' does not, and on top of that, taking the stable limit of cohomology of $\overline{\mathcal{M}}_{g,n}$ requires a finer tuning, cf. Pikaart [Pik95].

Our plan here is to show that a k -minimal model of $\mathcal{M}_{g,n}$ is independent of k as long as $g \geq 2k + 3$. In particular, the limit of k -minimal models exists and may be called a “ k -minimal model of $\mathcal{M}_{\infty,n}$ ”, continuing the abuse of notation adopted for cohomology. Since a minimal model of a space may be obtained as a union of k -minimal models, we call this union a “minimal model of $\mathcal{M}_{\infty,n}$ ”. Each of these k -minimal models is k -formal in a natural sense, see below, and the formality of $\mathcal{M}_{\infty,n}$ follows.

First of all, we recall basic notions on k -minimal models, see [GM81, Leh90, Mor78]. From now on we will assume that our spaces and algebras are connected and simply connected, i.e., their $H^0 = \mathbb{C}$ and $H^1 = 0$. In application to moduli spaces, this is the case as long as $g \geq 1$, see Harer [Har88]. A DG algebra $\mathfrak{M}$ is called *minimal* if it is free as a DG commutative algebra, $\mathfrak{M}^1 = 0$, and $d(\mathfrak{M}) \subset \mathfrak{M}^+ \cdot \mathfrak{M}^+$, where $\mathfrak{M}^+ = \bigoplus_{i>0} \mathfrak{M}^i$. A *minimal model* of a DG algebra $\mathcal{A}$ is a minimal DG algebra $\mathfrak{M}$ along with a *quasi-isomorphism* $\mathfrak{M} \rightarrow \mathcal{A}$, a morphism of DG algebras inducing an isomorphism on cohomology. Every DG algebra $\mathcal{A}$ has a minimal model, unique up to an isomorphism, which is in its turn unique up to homotopy.

Let $k \geq 0$ be an integer. A k -minimal model of a DG algebra $\mathcal{A}$ is a minimal algebra $\mathfrak{M}(k)$ generated by elements in degrees $\leq k$ along with a morphism $\mathfrak{M}(k) \rightarrow \mathcal{A}$ inducing an isomorphism on cohomology in degrees $\leq k$ and an injection in degree $k+1$. A k -minimal model is unique up to an isomorphism uniquely defined up to homotopy. If one has an increasing sequence of embeddings

$$\mathfrak{M}(0) \subset \mathfrak{M}(1) \subset \mathfrak{M}(2) \subset \dots$$

together with morphisms $\mathfrak{M}(k) \rightarrow \mathcal{A}$ compatible with each other, so that $\mathfrak{M}(k)$ is a k -minimal model of $\mathcal{A}$, then the union $\mathfrak{M} = \bigcup_k \mathfrak{M}(k)$ along with the natural morphism $\mathfrak{M} \rightarrow \mathcal{A}$ is a minimal model of $\mathcal{A}$.

We will call a DG algebra $\mathcal{A}$ (or a space whose rational homotopy type we consider) *formal* if a minimal model $\mathfrak{M}$ of $\mathcal{A}$ is isomorphic to a minimal model of its cohomology $H^\bullet(\mathcal{A})$ taken with the zero differential. This is equivalent to saying that there is a quasi-isomorphism $\mathfrak{M} \rightarrow H^\bullet(\mathcal{A})$. In this case the rational homotopy type of $\mathcal{A}$ (or the space) is determined by its rational cohomology ring. Formality implies that all Massey products are zero. If the cohomology algebra $H^\bullet(\mathcal{A})$ is free as a graded commutative algebra, then $\mathcal{A}$ is formal; see Proposition 3.1 for a k -version of this statement. Another example of a formal space is any compact Kähler manifold, a famous result of Griffiths, Deligne, Morgan, and Sullivan [DGMS75]. We will similarly call a DG algebra $\mathcal{A}$ or a space k -formal, if k -minimal models of $\mathcal{A}$ and its cohomology are isomorphic.

We also need to prove the following proposition establishing a particular case of k -formality. We say that a graded commutative algebra $\mathcal{C}$ is k -free if there exists a graded vector space $V = \bigoplus_{i=0}^k V^k$ and a mapping $V \rightarrow \mathcal{C}$ of graded vector spaces defining a morphism $S(V) \rightarrow \mathcal{C}$ of graded algebras, where $S(V)$ is the free DG commutative algebra on V , which is an isomorphism in degrees $\leq k$ and an injection in degree $k+1$.

PROPOSITION 3.1. *Suppose that the cohomology $H^\bullet(\mathcal{A})$ of a DG algebra $\mathcal{A}$ is k -free, based on a graded vector space V . Then $S(V)$ is a k -minimal model of $\mathcal{A}$, and $\mathcal{A}$ is k -formal.*

PROOF. If $H^\bullet(\mathcal{A})$ is k -free, then there exists a linear injection $\phi : V \hookrightarrow H^\bullet(\mathcal{A})$. Pick a linear mapping $V \rightarrow \mathcal{A}$ which takes each element v of V to a cocycle representing the cohomology class $\phi(v)$. Then the natural morphism of graded commutative algebras $S(V) \rightarrow \mathcal{A}$ obviously respects the differentials and satisfies the axioms of a k -minimal model of $\mathcal{A}$. By assumption, $S(V)$ is at the same time a k -minimal model of $H^\bullet(\mathcal{A})$, whence k -formality of $\mathcal{A}$. $\square$

Now we are ready to present the main result of the paper.

- THEOREM 3.2.**
1. *The moduli space $\mathcal{M}_{g,n}$ is k -formal for $g \geq 2k+3$.*
 2. *The subalgebra $H^\bullet(\mathcal{M}_{\infty,n})(k)$ of $H^\bullet(\mathcal{M}_{\infty,n})$ generated in degrees $\leq k$ is a k -minimal model of $\mathcal{M}_{g,n}$ for $g \geq 2k+3$.*
 3. *A k -minimal model of $\mathcal{M}_{g,n}$ is independent of g as long as $g \geq 2k+3$. We will call it a k -minimal model of $\mathcal{M}_{\infty,n}$.*
 4. *The k -minimal models of $\mathcal{M}_{\infty,n}$ form an increasing sequence of embeddings. The union, a minimal model of $\mathcal{M}_{\infty,n}$, is isomorphic to its cohomology $H^\bullet(\mathcal{M}_{\infty,n})$. In particular, $\mathcal{M}_{\infty,n}$ is formal.*

PROOF. 1. The stable cohomology $H^\bullet(\mathcal{M}_{\infty,n})$ is a free graded commutative algebra, *i.e.*, isomorphic to $S(V)$ for a graded vector space V , according to Miller-Morita's theorem [Mil86] for $n = 0$ and Looijenga's handling [Loo96] of the $n \geq 0$ case. Moreover, $H^0(\mathcal{M}_{\infty,n}) = \mathbb{C}$ and $H^1(\mathcal{M}_{\infty,n}) = 0$ for $g \geq 1$, see Harer [Har88]. Given a nonnegative integer k , the groups $H^k(\mathcal{M}_{g,n})$ are known to stabilize as soon as $g \geq 2k + 1$; this is the Harer-Ivanov Stability Theorem [Har85, Iva93]. Therefore, $H^\bullet(\mathcal{M}_{g,n})$ is k -free for $g \geq 2k + 3$: the mapping $V^{\leq k} \rightarrow H^\bullet(\mathcal{M}_{g,n})$ makes $H^\bullet(\mathcal{M}_{g,n})$ a k -free graded algebra. Proposition 3.1 then implies that $\mathcal{M}_{g,n}$ is k -formal for $g \geq 2k + 3$, $S(V^{\leq k})$ being a k -minimal model of it.

2 and 3. Notice that since $H^\bullet(\mathcal{M}_{\infty,n})$ is free, the subalgebra $H^\bullet(\mathcal{M}_{\infty,n})(k)$ generated in degrees $\leq k$ can be identified with $S(V^{\leq k})$, which we have just seen to be a k -minimal model of $\mathcal{M}_{g,n}$ for $g \geq 2k + 3$.

4. The subalgebras

$$H^\bullet(\mathcal{M}_{\infty,n})(0) \subset H^\bullet(\mathcal{M}_{\infty,n})(1) \subset H^\bullet(\mathcal{M}_{\infty,n})(2) \subset \dots$$

form an increasing sequence of k -minimal models of $\mathcal{M}_{\infty,n}$, therefore, their union, $H^\bullet(\mathcal{M}_{\infty,n})$, is a minimal model of $\mathcal{M}_{\infty,n}$, and thereby $\mathcal{M}_{\infty,n}$ is formal. $\square$

In view of this result, Mumford's Conjecture [Mum83], if true, implies the following refinement: *the polynomial algebra on the Mumford-Morita-Miller classes κ_i , $i = 1, 2, \dots$, and the first Chern classes $c_1(T_i)$ of the "tangent at the i th puncture" bundles, $i = 1, 2, \dots, n$, with a zero differential is the stable minimal model of the moduli space $\mathcal{M}_{g,n}$ as $g \rightarrow \infty$.*

References

- [Del71] P. Deligne. Théorie de Hodge. II. *Inst. Hautes Études Sci. Publ. Math.*, (40):5–57, 1971.
- [DGMS75] P. Deligne, P. A. Griffiths, J. W. Morgan, and D. Sullivan. Real homotopy theory of Kähler manifolds. *Invent. Math.*, 29(3):245–274, 1975.
- [GJ94] E. Getzler and J. D. S. Jones. Operads, homotopy algebra and iterated integrals for double loop spaces. Preprint, Department of Mathematics, MIT, March 1994. [hep-th/9403055](#).
- [GM81] P. A. Griffiths and J. W. Morgan. *Rational homotopy theory and differential forms*, volume 16 of *Progress in Mathematics*. Birkhäuser, Boston, MA, 1981.
- [Har85] J. L. Harer. Stability of the homology of the mapping class groups of orientable surfaces. *Ann. of Math. (2)*, 121(2):215–249, 1985.
- [Har88] J. L. Harer. The cohomology of the moduli space of curves. In *Theory of moduli (Montecatini Terme, 1985)*, volume 1337 of *Lecture Notes in Math.*, pages 138–221. Springer, Berlin, 1988.
- [HS79] S. Halperin and J. Stasheff. Obstructions to homotopy equivalences. *Adv. in Math.*, 32(3):233–279, 1979.
- [Iva93] Nikolai V. Ivanov. On the homology stability for Teichmüller modular groups: closed surfaces and twisted coefficients. In *Mapping class groups and moduli spaces of Riemann surfaces (Göttingen, 1991/Seattle, WA, 1991)*, volume 150 of *Contemp. Math.*, pages 149–194. Amer. Math. Soc., Providence, RI, 1993.
- [Kon94] M. Kontsevich. Feynman diagrams and low-dimensional topology. In *First European Congress of Mathematics, Vol. II (Paris, 1992)*, volume 120 of *Progr. Math.*, pages 97–121. Birkhäuser, Basel, 1994.
- [KSV95] T. Kimura, J. Stasheff, and A. A. Voronov. On operad structures of moduli spaces and string theory. *Commun. Math. Phys.*, 171:1–25, 1995. [hep-th/9307114](#).
- [KSV96] T. Kimura, J. Stasheff, and A. A. Voronov. Homology of moduli of curves and commutative homotopy algebras. In *The Gelfand Mathematical Seminars, 1993–1995*, Gelfand Math. Sem., pages 151–170. Birkhäuser Boston, Boston, MA, 1996. [alg-geom/9502006](#).

- [KVZ97] T. Kimura, A. A. Voronov, and G. J. Zuckerman. Homotopy Gerstenhaber algebras and topological field theory. In *Operads: Proceedings of Renaissance Conferences (Hartford, CT/Luminy, 1995)*, volume 202 of *Contemp. Math.*, pages 305–333, Providence, RI, 1997. Amer. Math. Soc. [q-alg/9602009](#).
- [Leh90] D. Lehmann. Théorie homotopique des formes différentielles (d’après D. Sullivan). *Astérisque*, (45):145, 1990.
- [Loo96] E. Looijenga. Stable cohomology of the mapping class group with symplectic coefficients and of the universal Abel-Jacobi map. *J. Algebraic Geom.*, 5(1):135–150, 1996.
- [Mil86] E. Y. Miller. The homology of the mapping class group. *J. Differential Geom.*, 24(1):1–14, 1986.
- [Mor78] J. W. Morgan. The algebraic topology of smooth algebraic varieties. *Inst. Hautes Études Sci. Publ. Math.*, (48):137–204, 1978.
- [Mum83] D. Mumford. Towards an enumerative geometry of the moduli space of curves. In *Arithmetic and geometry, Vol. II*, volume 36 of *Progr. Math.*, pages 271–328. Birkhäuser Boston, Boston, Mass., 1983.
- [Pik95] M. Pikaart. An orbifold partition of $\overline{M}_g^n$. In *The moduli space of curves (Texel Island, 1994)*, volume 129 of *Progr. Math.*, pages 467–482. Birkhäuser Boston, Boston, MA, 1995.
- [Sta63] J. Stasheff. Homotopy associativity of H -spaces. I, II. *Trans. Amer. Math. Soc.*, 108:275–312, 1963.
- [Sul77] D. Sullivan. Infinitesimal computations in topology. *Inst. Hautes Études Sci. Publ. Math.*, (47):269–331, 1977.

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, SAKYO-KU, KYOTO, 606-01, JAPAN AND DEPARTMENT OF MATHEMATICS, M.I.T., 2-270, 77 MASSACHUSETTS AVE., CAMBRIDGE, MA 02139-4307, USA

E-mail address: voronov@kurims.kyoto-u.ac.jp